

REVIEW THE INFORMATION IN THIS DOCUMENT CAREFULLY.
IF YOU DO NOT UNDERSTAND ANY PART OF THIS DOCUMENT,
ASK YOUR HEALTH CARE PROVIDER FOR HELP.

IF YOU ARE HAVING PROBLEMS WITH YOUR SPINAL CORD,
YOUR PHYSICIAN WILL WANT TO KNOW ABOUT YOUR
PAIN AND HOW IT AFFECTS YOUR LIFE. YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

Musculoskeletal pain

IF YOU HAVE MUSCULOSKELETAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE MUSCULOSKELETAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE MUSCULOSKELETAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE MUSCULOSKELETAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE MUSCULOSKELETAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.



Visceral pain

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

Managing pain after SCI

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

Physical therapy

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

IF YOU HAVE VISCERAL PAIN, YOUR PHYSICIAN
WILL WANT TO KNOW ABOUT YOUR HISTORY OF PAIN,
YOUR CURRENT PAIN, AND HOW YOUR PAIN AFFECTS
YOUR LIFE.

- **Transcutaneous electrical nerve stimulation (TENS)** is sometimes used to treat musculoskeletal pain. Electrodes are placed on the surface of the skin and send low levels of electrical current into the body. The current blocks signals from the areas of nerve damage that are triggering a pain response.

Behavioral psychological treatments for pain

We now know that people can learn to use psychological techniques to help them manage their pain better so it doesn't take over their lives. Psychologists trained in pain management can help with a variety of behavioral techniques proven to be effective in reducing the intensity and impact of pain.

- **Relaxation techniques and/or biofeedback** designed to teach you how to reduce muscle pain tension and "mental tension" associated with pain can be helpful in self-management.
- **Self-hypnosis training** has proven helpful for reducing chronic pain in some individuals.
- **Cognitive restructuring.** Learning how to think differently about your pain and its effects can lead to changes in brain activity and, in turn, the experience of pain.
- **Individual psychotherapy** designed to help identify desired goals and increase pleasure and meaning in daily life can help reduce pain. Therapy can also help if there is a significant amount of anxiety associated with pain.

Medications

There are many medications to treat pain. All of the medications listed below have shown some success in reducing pain, but none do so completely in every instance. All have possible side effects, some of which can be serious. Discuss all side effects with your doctor. Sometimes combinations of drugs work better than a single drug.

- **Non-steroidal anti-inflammatory drugs** (also known as NSAIDs) such as aspirin, ibuprofen (Motrin, Advil) and naproxen are commonly used to treat musculoskeletal pain. Side effects may include stomach upset or bleeding.
- **Antiseizure medications** such as gabapentin (Neurontin) and Pregabalin (Lyrica) are used to treat neuropathic pain. Side effects include dizziness, sleepiness and swelling.

- **Antidepressants** are used to treat neuropathic pain and depression. These medications include selective serotonin norepinephrine reuptake inhibitors (SSNRIs), such as venlafaxine (Effexor), and tricyclics, such as amitriptyline (Elavil). Side effects include dry mouth, sleepiness, dizziness, and (with SSNRIs) nausea.
- **Narcotics (opiates)** such as morphine, codeine, hydrocodone and oxycodone are not recommended for treatment of chronic or long standing neuropathic or musculoskeletal pain. These drugs have many side effects, including constipation, depression of breathing, and slowed thinking. They are habit forming and use often leads to dependency. Withdrawal symptoms can occur when stopped suddenly.
- **Muscle relaxants and anti-spasticity** medications such as diazepam (Valium), baclofen (Lioresal) and tizanidine (Zanaflex) are used to treat spasm-related and musculoskeletal pain. These may be taken by mouth or delivered directly to the spinal cord through an implanted pump (see "Intrathecal pumps" below). These drugs can cause sleepiness, confusion and other side effects.
- **Topical local anesthetics** such as lidocaine (Lidoderm) are used to treat pain that occurs when skin is touched lightly (called allodynia).

Surgical Treatments

- **Dorsal column stimulator** is used to treat neuropathic pain due to nerve root damage. A high frequency, low intensity nerve stimulator is surgically placed in the spinal canal next to the spinal cord or nerve roots.
- **Intrathecal pumps** are used to treat neuropathic pain (using morphine) or muscle spasm-related pain (using baclofen) A pump containing morphine or baclofen is surgically placed under the skin in the abdomen. It delivers the medication directly to the spinal cord and nerve roots.

Prevention and self-care

- **Get treatment for medical problems.** Overall health can have a big effect on pain. Urinary tract infections, bowel problems, skin problems, sleep problems and spasticity can make pain worse or harder to treat. Keeping yourself as healthy as possible can help reduce pain.
- **Try to get as much exercise as possible.** Getting regular physical activity can reduce pain as well as improve mood and overall health. It can also be enjoyable and distract you from pain. Your health provider can help you choose physical activities that are safe and appropriate for you. Also see the supplement “**Activity Modification for Musculoskeletal Pain.**”
- **Get treatment for depression.** Depression can make pain worse. It is best treated through counseling and medication. Getting treatment for depression can help you cope with chronic pain and improve your quality of life.
- **Reduce stress** Stress can make pain worse or make the pain harder to manage. You can learn to manage stress through counseling and learning techniques to reduce stress and tension, such as relaxation training, biofeedback and hypnosis. Exercise helps reduce stress.
- **Distract yourself.** Distraction is one of the best methods for coping with chronic pain. Participating in enjoyable and meaningful activities can help reduce pain and help you feel more in control of your life, especially when pain is at its worst. When we are bored and inactive, we tend to focus more on pain, and this can make pain feel worse.
- **Keep a record.** Everyone’s pain is a little different. Keep a record of what makes you feel better and what makes pain worse. Understanding things that affect your pain will help you and your doctor find effective ways to reduce pain.
- **Get a wheelchair seating evaluation.** Poor posture and improper seating can cause serious pain problems. Get your seating evaluated by a physical therapist who specializes in wheelchair seating. If you use a manual wheelchair, try to get

a high-strength, fully customizable chair made of the lightest material possible (aluminum or titanium). Learn the proper wheelchair propulsion (pushing) technique from a physical therapist.

- **Don't drink to ease pain.** Using alcohol as a pain medication can lead to alcohol abuse and other serious problems. Some medications should not be mixed with alcohol. Ask your doctor about drinking alcohol, and always read the labels of your prescriptions.

Finding help

It is important to get treatment for pain. The ideal source of help would be a physician and psychologist familiar with SCI and pain management, working together.

If you do not have access to such experts, the next best alternative is to seek help from a multidisciplinary pain clinic where physicians and psychologists are available. Work closely with a health care provider with whom you trust and who understands your condition.

Chronic pain is not hopeless. Try not to become discouraged if one treatment doesn’t work, and be open to trying different techniques. While complete relief from pain may not be possible, living better despite pain is a realistic goal.

Resources

- Pain Connection, www.painconnection.org
- American Pain Society, www.ampainsoc.org
- American Pain Foundation, www.painfoundation.org
- CareCure Community Moderated Forums, including a pain forum. <http://sci.rutgers.edu/forum/>

Source

Our health information content is based on research evidence and/or professional consensus and has been reviewed and approved by an editorial team of experts from the SCI Model Systems.

Authorship

Pain after Spinal Cord Injury was developed by J. Scott Richards, PhD, Trevor Dyson-Hudson, MD, Thomas N. Bryce, MD, and Anthony Chiodo, MD, in collaboration with the University of Washington Model Systems Knowledge Translation Center.

Portions of this document were adapted from materials developed by the UAB SCI Model System, University of Michigan Model SCI Care System, Northwest Regional SCI System, and Craig Hospital.